

Karyological study of the Caspian bent-toed Gecko *Cyrtopodion caspium* (Sauria: Gekkonidae) from North and North-Eastern of Iran

Farahnaz Molavi^{1*}, Haji-Gholi Kami², Morteza Yazdanpanahi¹

1-Department of Biology, Faculty of sciences, Mashhad Branch, Islamic Azad University, Mashhad, Iran

2-Department of Biology, Faculty of Sciences, Golestan University, Gorgan, Iran

Received 21 January 2014

Accepted 01 March 2014

Abstract

The diploid number among Gekkonid lizards ranges from $2n=16$ to $2n=46$. The majority of geckos' chromosomes are small which makes the study of their chromosomes very difficult. For this reason, karyotype of *Cyrtopodion caspium* and some similar gecko are still not known. The close relatives of *C. caspium* have variation in the chromosomal number in their populations. So to find out possible variability among these populations in Iran. We described the karyotype of testis and bone marrow of 14 specimens of this species from seven provinces and 10 localities in north and north-eastern parts of Iran in 2011. They showed $2n=38$ for all populations. There is not intraspecific chromosome polymorphism in *Cyrtopodion caspium*. The karyotype consists of one pair of metacentric element and 17 pairs of telocentric or subtelocentric elements and one pair of three arm chromosomes. The number of haploid chromosomes was 21 for three male specimens. The chromatid separation did not show heteromorphic sex chromosomes. Our results highlight the need for continued research into the basic biology and taxonomy of *Cyrtopodion caspium* in Iran.

Keywords: Chromosome structure, Karyology, Sex chromosome, Lizard, Iran, and Caspian bent toed gecko

Introduction

The family Gekkonidae is the most diverse and the oldest group of reptiles which have a worldwide distribution (Anderson, 1999). It is the largest family of lizards, comprising 100 genera and 943 species. Although there are several studies on this species, a few of them are related to the karyotype of this family (Ahmadzadeh *et al.*, 2004, 2005, 2008; Hojati *et al.*, 2009).

Cyrtopodion caspium is a small sized oviparous nocturnal lizard, as a house gecko, it is distributed in many cities and villages in Iran. It consists of two subspecies, *Cyrtopodion caspium caspium* (Eichwald, 1831) and *Cyrtopodion caspium insularis* (Akhmedov and Szczerbak, 1978). The main habitats of *C. c. caspium* in Iran are located in the Gorgan region of Mazandaran, to northern and eastern Khorasan, extending south to Sistan and west to Azerbaijan. *C. c. insularis* has only been reported from a limited region in the Caspian Sea. (Rastegar-Pouyani *et al.*, 2008; Rhodin *et al.*, 2010).

The diploid number among Gekkonid lizards ranges from $2n=16$ to $2n=46$ (Schmidt *et al.*,

1994). The typical karyotype consists of a gradual series of acrocentric chromosomes which there is no difference between macro and micro chromosomes.

The sex determination mechanisms in Saurian have not been completely understood. Sex chromosome evolution in recent data from some lizard's families suggests that they have multiple origins (Beak, 1983). There are species with chromosomal sex determination mechanisms ascribed to male heterogamete in the family's Iguanidae (Frost & Etheridge, 1989), Lacertidae, Teiidae, Scincidae, and Pygopodidae. Female heterogamety is known in the family's Gekkonidae, Varanidae, and Lacertidae (Peccinini-Seale *et al.*, 1981). In the genus *Cnemidophorus*, Cole *et al.* (1969) and Bull (1978) reported a chromosomal sex determination mechanism of the type XX: XY for *Cnemidophorus tigris*. We performed a comparative karyological analysis among different populations of *Cyrtopodion caspium*, from North and North-Eastern Iran, in order to find out possible variability among these populations.

*Corresponding author E-mail:
fm_yazdan@yahoo.com

Material and Methods

During the field work from May to June 2011, we have collected 14 adult specimens (9 males and 5 females) of *Cyrtopodion caspium* from seven provinces and 10 localities of north and north-eastern of Iran (Table 1). All individuals of these species were transferred to the laboratory at the IAU branch of Mashhad, and their karyotypes were determined from all specimens.

Chromosomal preparations were obtained from suspension of bone marrow cells from femur and vertebral column and of spleen cells according to the procedures of Porter and Sites (1986). Meiotic chromosomal spreads were prepared from testis suspension adapted for lizards by Peccinini-Seale *et al.* (1971). Metaphase plates were obtained by the squash method, with treatment of vinblastine specimens. For 9 male individuals at least 11-21 complete metaphases were analyzed. Slides were stained by 4% gimsa with pH 7.2. The photographs were taken on a Zeiss microscope with 100X magnification. Chromosome number and properties were determined by MIP Software.

Table 1. Sampling localities of *Cyrtopodion caspium*

	Locality- province	Longitude	Latitude	Altitude (m)
1	Anzali port- Gilan	49° 25'	37° 28'	15
2	Siyahdarreh- Gilan	49° 08'	37° 07'	10
3	Fereydounkenar- Mazandaran	53° 23'	36° 50'	-10
4	Zaghmarz- Mazandaran	53° 33'	36° 43'	-10
5	Ziyarat village-Golestan	54° 24'	36° 54'	250
6	Alagol wetland- Golestan	54° 36'	37° 17'	00
7	Kalat- Northern Khorasan	59° 40'	37° 00'	1000
8	Ahmad Abad village - Semnan	56° 42'	35° 48'	1000
9	Shandiz - Razavi Khorosan	59° 25'	33° 56'	1360
10	Byrjand - Southern Khorasan	59° 13'	32° 53'	1491




Figure 1. The map of sampling localities.

Results

The karyotype of *Cyrtopodion caspium* is shown in figure 2. Metaphase analyses showed a diploid number of 38 chromosomes ($2n=38$). The fifth pair was metacentric and other pairs were subtelocentric or telocentric, and there is a pair with three chromatids. Macro and microchromosomes were indistinguishable and no sex chromosome polymorphism could be observed (Figure 1 and Table 2 and 3).




Figure 2. Karyotype of *Cyrtopodion caspium*.

Table 3. Calculations on karyotype of *Cyrtopodion caspium*.

Order	Variable name	Value
1	2n	38
2	Number of Arms	46
3	Fn	44
4	Fna	2
5	All Chromosome Length	1490.08 μ m
6	Total Metacentric	0
7	Total Submetacentric	8
8	Total Acrocentric	24
9	Total Telocentric	6

No heteromorphic sex chromosomes were observed in the karyotype of *Cyrtopodion caspium*. Meiotic analysis showed a haploid number of 21 chromosomes (n=21) for male specimens (Figure 3).




Figure 3. Karyotype of spermatid cell in *C. caspium*.

There was variability in number (1-2 nucleoli) and size (small, medium and large) of nucleoli in the 50 analyzed interphase cells (Figure 4).




Figure 4. Interphasic nuclei of bone marrow cells.

However, T.F-factor that stands for symmetry is lower than 50% and shows that chromosomes are not in symmetrical position and are located very far from chromosomal symmetry.

$$T.F\% = \frac{\text{Short arms lengths}}{\text{Total arms lengths}} \times 100$$

T.F. = 1.594

Moreover, to show the symmetrical position the DRL index is employed which is a maximum relative length of chromosomes minus from minimum relative length of chromosomes. DRL

index equals 0.1069 that confirms lack of symmetrical condition in the species.

Table 2. Description of *Cyrtopodion caspium* chromosomes.

No.	Length (µm)	Up Arm (µm)	Down Arm (µm)	Lcn/L total	Centromer Index (µm)	Chromosome Type
1	93.66	24.6	69.07	0.063	0.26	Acrocentric
2	76.48	29.07	47.41	0.051	0.38	Acrocentric
3	72.47	33.06	39.41	0.049	0.46	Sub-metacentric
4	74.09	25.08	49.01	0.05	0.34	Sub-metacentric
5	83.41	5.39	78.03	0.056	0.06	Telocentric
6	80.35	6.32	74.03	0.054	0.08	Telocentric
7	50.18	5	45.18	0.034	0.1	Acrocentric
8	50.43	5.83	44.6	0.034	0.12	Acrocentric
9	43.05	5.1	37.95	0.029	0.12	Telocentric
10	47.01	5	42.01	0.032	0.11	Telocentric
11	35.45	12.04	23.41	0.024	0.34	Submetacentric
12	18.3	8.25	10.05	0.012	0.45	Submetacentric
13	40.54	5.1	35.44	0.027	0.13	Telocentric
14	41	5.1	35.9	0.028	0.12	Telocentric
15	34.96	10.44	24.52	0.024	0.3	Submetacentric
16	37.25	12.17	25.08	0.025	0.33	Submetacentric
17	33.14	9.06	24.08	0.022	0.27	Acrocentric
18	35.06	12.04	23.02	0.024	0.34	Acrocentric
19	31	5	26	0.021	0.16	Acrocentric
20	39.11	5.1	34.01	0.026	0.13	Acrocentric
21	26.49	9.49	17	0.018	0.36	Submetacentric
22	32.7	7.07	25.63	0.022	0.22	Submetacentric
23	26	5	21	0.017	0.19	Acrocentric
24	24.41	5.1	19.31	0.016	0.21	Acrocentric

25	22.36	6.08	16.28	0.015	0.27	Acrocentric
26	22.19	5.1	17.09	0.015	0.23	Acrocentric
27	21	5	16	0.014	0.24	Acrocentric
28	20.96	5.83	15.13	0.014	0.28	Acrocentric
29	18.57	6.4	12.17	0.012	0.34	Acrocentric
30	19.66	8	11.66	0.013	0.41	Acrocentric
31	18.73	5.39	13.34	0.013	0.29	Acrocentric
32	19	5	14	0.013	0.26	Acrocentric
33	18.34	5	13.34	0.012	0.27	Acrocentric
34	16.67	7.62	9.06	0.011	0.46	Acrocentric
35	12.22	3	9.22	0.008	0.25	Acrocentric
36	9.06	0	9.06	0.006	0	Acrocentric
37	59.74	5	54.74	0.04	0.08	Acrocentric
38	85.04	7.81	77.23	0.057	0.09	Acrocentric

Discussion

Chromosomal evolution has been occurred in all vertebrates. The results of other survey reveal that many minor chromosome rearrangements have been occurred. However, minor deletions throughout the genome of the ancestors of reptiles and birds have led to reduce birds' genome to 50% of reptiles' genome (Swanson *et al.* 1981; Edwards 2009; Iturra *et al.*, 1994).

Single or multiple centric fissions are the main chromosome rearrangement found in the evolution of lizard karyotype, including *Anolis*, *Sceloporus grammicus* and *Liolaemus* (Webster *et al.*, 1972; Lambrot and Alvares-Sarret, 1989; Sites, 1983; Lambrot, 1991)

There is not intraspecific chromosome polymorphism in *Cyrtopodion caspium*, which have not shown considerable chromosome variations in several examined populations. Although Intra-specific chromosomal polymorphism is generally between 10-50 % in lizards (Iturra *et al.*, 1994).

Some authors have argued that the $2n=38$ is ancestral for the family Gekkonidae (and perhaps the entire suborder), although others have expressed different opinions (Shibaïke *et al.*, 2009; Oliver *et al.*, 2007).

However all specimens from seven provinces were monomorphic. The diploid number of $2n=38$ found in *Cyrtopodion caspium* is situated within the

range of the family.

Different specimens of *Diplodactylus tessellatus* from Australia exhibit $2n=28$, 30, and 38 that $2n=38$ is the ancestral case. 38 chromosomes are reduced to 30 and 28 through minor fissions of some acrocentric chromosomes. Moreover, $2n=38$, 36, 34, 30 are reported from *Vittatus diplodactylus* (Oliver *et al.*, 2007).

In *Tarento lamauritanica* (Gekkonidae) only one chromosome number has been reported ($2n=42$). Many herpetologists believe that more studies need to investigate variation in chromosome number (Shibaïke *et al.*, 2009 and Hidestonhiet *al.* 2001). For example, many studies on *Diplodactylus vittatus* show high variation in chromosome number in members of Gekkonidae. The above species exhibits 3 different chromosome numbers from different regions. (King 1977) Geckoes' karyotypes are $2n=38$ in many cases. However, diversity in chromosome numbers has been observed in geckos. It should be emphasize that different chromosome numbers are available in populations which are similar morphologically. (Shibaïke *et al.*, 2009).

Gekko gekko (one of the near relatives of *C. caspium*) exhibits only $2n=38$ and there is no variation in chromosome number. However, it should be considered that only one specimen of *G. gekko* has been investigated to study the karyotype of this species. (Cohen *et al.* 1967)

Evolution of karyotypes occurs in neighboring geographical regions; for example, chromosome evolution of 9 Gekkonidae species are reported from Eastern Asian islands (*G. shibatai*, *G. tawaensis*, *G. vertebral* is, *G. yakuensis*, and 3 non-described species). Moreover the number of chromosomes are $2n=38$ in all of specimens. However different karyotypes have been investigated from different populations (Shitake and Takahashi, 2009).

There are many acrocentric chromosomes in all of reptiles (Hidetoshi *et al.*, 2001). The majority of geckoes' chromosomes are small which makes the study of their chromosomes very difficult (Katia and Machado, 1995).

Karyotypes with no distinction between macro and micro chromosomes, as the one we found in *C. caspium* are also typical of the Gekkonidae.

Heteromorphic sex chromosomes were not observed in *C. caspium*. It was corroborated by the absence of a distinctive heteromorphic bivalent in male cells. The presence of heteromorphic sex chromosome in Gekkonidae is known only from 10 species belonging to five genera (Shitake and Takahashi, 2009). However as far as we know; there is no reference between sexual chromosomes in

Cyrtopodion. (Katia and Machado, 1995)

Most Gekkonids have a chromosomal mechanism of sex determination of the ZZ:ZW type (Kawai *et al.*, 2009), in which the heterogametic sex is the female, such as in *Heteronotia binoei* (Mortiz, 1990), *Gehyra australis* (King 1983), *Gehyra purpuracens* (Moritz, 1984) and *Cyrtodactylus pudisulcus* (Shitake and Takahashi 2009). But, in a few species, the mechanism is of the XX:XY type, such as in *Gekko gecko* (Solleder and Schmid 1984), *Gekko japonica* (Mortis 1990) and *Gonato desceciliae* (McBee *et al.*, 1987). *G. hokouensis* demonstrates high diversity of sex chromosomes and their rapid evolution; however, mechanisms of sex determination are evolved rapidly (Ezaz *et al.*, 2009).

Temperature is the most important factor in determination of sex in lizards, for example ancestral Lophosaurus (Janes *et al.*, 2009) but in some cases sex chromosomes determine gender (Janes *et al.*, 2009). For example, there are sex chromosomes in *Gekko hokouensis* and sex determination is not related to environmental factors in these species (Kawai *et al.*, 2009).

Acknowledgments

We appreciate the efforts of Narges Tehrani, who assisted in the collection of the lizards. Special thanks also to Dr. Darvish, because he was allowed to use the MIP software. This software is limited to Rodentology Research Department (RRD).

References

1. Ahmadzadeh F. (2004) A preliminary study of the Lizard fauna and their habitats in Meshkinshahr district. *Environmental Sciences*. 1(2). 39-44.
2. Ahmadzadeh F., Kiabi B. H., Kami H. G., and V. Hojati. (2008) A preliminary study of the Lizard fauna and their habitats in Northwestern Iran. *Asiatic Herpetological Research*. Vol 11, pp. 1-9.
3. Ahmadzadeh F., Kiabi B. H., Zehzad B., and Kami H. G. (2005) A survey on Lizard fauna in the northern part of Ardebil province of Iran. First international conference of biology. 23-25 August. Gilan University. Rasht. Iran.
4. Anderson S. C. (1999) The lizards of Iran. *Contributions to Herpetology*. Vol. 15, Society for the Study of Amphibians and Reptiles, Saint Louis, Missouri: i-vii, 35-44.
5. Beçak W. (1983) Evolution and differentiation of sex chromosome in lower vertebrates. *Differentiation*, 23(supl.): 3-12.
6. Bull J. J. (1978) Sex chromosome differentiation: an intermediate stage in a lizard. *Canadian Journal of Genetics and Cytology*, 20: 205-209.
7. Cohen M. M., Huang Ch., and Clrak H. F.. (1967) The somatic chromosomes of 3 lizard species: *Gekko gecko*, *Iguana iguana*, and *Crotaphytus collaris* cellular and molecular life sciences. Vol. 23, No. 9, 769-771
8. Cole C. J., Lowe C. H., and Wright J. W. (1969) Sex Chromosomes in Teiid Whiptail Lizards (Genus *Cnemidophorus*). *American Museum Novitates*, 2395: 1-14.
9. Eric J. V., and Bruce T. L. (2006) Multiple independent origins of sex chromosomes in amniotes PNAS. 103(20): 18031–18048
10. Ezaz T., Quinn A. E., Sarre S. D., O'Meally D., Georges A., Marshall J. A., and Graves M.. (2009) Molecular marker suggests rapid changes of sex-determining mechanisms in Australian dragon lizards, *Chromosome Research*, Vol.17, 91-98.
11. Frost D. R., and Etheridge R. (1989) A phylogenetic analysis and taxonomy of iguanian lizards (Reptilia: Squamata). *Misc. Publ. Mus. Nat. Hist. Univ. Kansas*, 81: 1-65.
12. Hidetoshi O., Hikida T., Nabhitabhata J., and Panha S.. (2001) Cryptic Taxonomic Diversity in Two Broadly Distributed Lizards of Thailand (*Mabuya macularia* and *Dixonius siamensis*) as Revealed by Chromosomal Investigations (Reptilia: Lacertilia) *The Natural History Journal of Chulalongkorn University*. 1(1): 1-7.
13. Hojati V., Kami H. G., Faghiri A., and Ahmadzaheh F. (2009) A morphological study of Bedriaga plate-tailed *Gecko*, *Teratoscincus bedriaga* Nikolsky 1900, in Semnan province of Iran (Reptilia: Gekkonidae), *Zoology in the Middle East*. 46: 113-115.
14. Iturra P., Veloso A., Espejo P., and Navarro J. (1994) Karyotypic and meiotic evidence for a Robertsonian chromosome polymorphism in the lizard *Liolaemus fuscus* (Tropiduridae, Sauria). *Brazilia Journal of Genetics*. 17(2): 171-174.
15. Janes, D. E., Ezaz T., Graves J. A. M., and Edwards S. V. (2009) Recombination and nucleotide diversity in the pseudoautosomal region of minimally

- differentiated sex chromosomes in the Emu, *Dromaius novaehollandiae*. Journal of Heredity 100(2): 125-136.
16. Katia C., and Machado P. (1995) C-and R-banding patterns and nucleolus organizer regions in the karyotype of *Hemidactylus mabouia* (Sauria, Gekkonidae). Brazilia Journal of Genetics. 18, p. 527-531.
 17. Kawai A., Junko I., Nishida C., Ayumi K., and Ota H.. (2009) The ZW sex chromosomes of *Gekko hokouensis*(Gekkonidae, Squamata) represent highly conserved homology with those of avian species. Chromosoma 118: 51-43.
 18. King M. (1977) Chromosomal and morphometric variation in the *GekkoDiplodactylus vittatus*. Austerallian Journal of Zoology 25(1): 57-43
 19. King M. (1983) Katyotypic evolution in *Gehyra* (Gekkonidae: Reptilia). III. The *Gehyra australis* complex. Aust. J. zool. 31: 723-741.
 20. Klug M. (1991) Checklist of gekkonoid lizards. Smithsonian Herpetological Information Service 85: 1-35.
 21. Lamborot M., and Alvares-Sarret E. (1989) Karyotypic characterization of some *Liolaemus* lizards in Chile (Iguanidae). Genome. 32: 393-403.
 22. Lamborot, M. (1991) Karyotypic variation among populations of *Liolaemus monticola* (Tropiduridae) separated by riverine barriers in the Andean Range. Copeia (4): 1044-1059.
 23. Levan A., Fredga K., and Sandberg A. A. (1964) Nomenclature for centromeric position on chromosomes. Hereditas, 52(2): 201-220.
 24. McBee K., Bickham J. W., and Dixon J. R. (1987) Male heterogamy and chromosomal variation in Caribbean geckos. Journal of Herpetology 21 (1): 68-71.
 25. Mortiz C. (1984) The evolution of a highly variable sex chromosome in *Gehyra purpurascens* (Gekkonidae): Chromosome 90: 111-119.
 26. Mortiz C. (1990) Patterns and processes of sex chromosome evolution in Gekkonid lizards (Reptilia: Sauria). In Cytogenetics of Amphibians and Reptiles (Olmo, E., Ed.). BirkhauserVerlag, Switzerland, pp. 205-220.
 27. Oliver P. M., Hutchinson M. and Hutchinson R. (2007) Karyotypic variation in the Australian gecko *Ciplodactylus tessellatus*, with the description of a new karyotypic complement for diplodactyline geckos Journal of Herpetology 41 (3): 540-543.
 28. Peccinini-Seale D. (1981) New developments in vertebrate cytotaxonomy. IV. Cytogenetic studies in reptiles. Genetica, 56: 123-148.
 29. Peccinini-Seale D., and Frota-Pessoa O. (1974) Structural heterozygozity in parthenogenetic populations of *Cnemidophorus lemniscatus* (Sauria: Teiidae) from the Amazonas Valley. Chromosome, 47: 439-451.
 30. Porter C. A., and Sites J. W. (1986) Evolution of *Sceloporus grammicus* complex (Sauria: Iguanidae) in central Mexico: population cytogenetics. Syst. Zool. 35: 334-358.
 31. Rastegar-Pouyani N., Kami H. G., Rajabzadeh M., Shafiei S., and Anderson S. C. (2008) Annotated check list of amphibians and reptiles of Iran. Iranian Journal of Animal Biosystematics(IJAB). 4 (1), 43-66.
 32. Rhodin A. G. J., Dijk P. P. V., Iverson J. B., Shaffer H. B. (2010) Turtle of the world: Annotated check list of taxonomy, synonymy, distribution and conservation Status. Chelonian Research Monographs. (5). 85-164.
 33. Schmid M., Feichtinger W., Nanda I., Schakowski R., Garcia R. V., Puppo J. M., and Badillo A. F. (1994) An extraordinary low diploid chromosome number in the reptile *Gonatodestaniae* (Squamata, Gekkonidae). J. Heredity. 85: 255-260.
 34. Scott C. C., and Edwards V. (2009) Genome evolution in Reptilia: in silico chicken mapping of 12,000 BAC-endsequences from two reptiles and a basal bird, Proceedings of the Avian Genomics Conference and Gene Ontology Annotation Workshop.
 35. Shibaïke, Y., Takahashi Y., Arikura I., Iizumi R., Kitakawa S., Sakai M., Imaoka C., Shiro H., Tanaka H., Akakubo N., Nakano M., Watanabe M., Ohne K., Kubota S., Kohno S., and Ota H. (2009) Chromosome evolution in the lizard genus *Gekko* (Gekkonidae, Squamata, Reptilia) in the East Asian islands. Cytogenetic and Genome Research.127:182-190.
 36. Sites J. W. (1983) Chromosome evolution in the iguanid lizard *Sceloporus*

- grammicus*. Chromosome polymorphisms. *Evolution* 37: 38-53.
37. Swanson C. P., Merz T., and William J. A. Y. (1981) *Cytogenetics: The chromosome in division, Inheritance and Evolution*. 2nd ed. Englewood Cliffs, N.J.: Prentice-Hall. xiv, 577pp.
 38. Webster T. P., Hall W. P., and Williams E. E. (1972) Fission in the evolution of a lizard karyotype. *Science*. 77: 611-613.