

Comparative Analysis of Commercial CCL21 and CCL21/IL1 β Recombinant Proteins by *in silico* Tools

Ahdiyeh Shahtaghi^{1†}, Ali Alam Shahnabadi^{1†}, Kamelia Kohannezhad¹, Neda Amini¹, Maria Beihaghi^{1,2*}

¹Department of Biology, Kavian Institute of Higher Education, Mashhad, Iran

²Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran

Received 11 March 2020

Accepted 16 May 2020

Abstract

One of the newest diagnostic methods and treatment of cancer is to design new drugs. It is now possible to design a drug with desired properties in theory and evaluate its therapeutic effects through bioinformatics tools. Among the studied drugs, those based on cytokine genes, which increase the body's immunity against cancer, are of great interest. Cytokines are small proteins that play an essential role in cell signaling and can affect the function and behavior of surrounding cells. CCL21 chemokine is one of the cytokines that possess antitumor properties has the potential for chemoattraction of T lymphocytes and dendritic cells. Interleukin 1 beta (IL1 β) is a cytokine involving different cellular activities such as the activation of neutrophils, B-Cells, and T-Cells. In the present study, we designed a drug-based cytokine gene to activate T cells and B cells by inserting defined CCL21 epitope and IL1 β peptide sequences into a protein construct. Molecular dynamics simulation was performed in Linux space using Gromex software. Results of RMSD, RMSF, and the radius of gyration obtained from the simulation showed the stability of both proteins, which indicated that there are no significant conformational differences between the commercial CCL21 and recombinant form. The interaction of synthetic construct and human CCL21 with the CCR7 receptor was also investigated by HADDOCK software. Obtained results showed no differences between these proteins, and recombinant protein has the same structural and conformational characteristics as human commercial CCL21.

Keywords: Cytokine, Chemokine, CCL21, Docking, Molecular Dynamics Simulation

Introduction

Nowadays, immunotherapy is a well-known method for understanding the problems related to the side effects of chemical drugs, analyzing the functional immune system during treatment, thus preventing tumor production. Among the drugs being studied for immunotherapy are cytokine-based drugs that increase immunity against cancer. Cytokines are small proteins that play an important role in cell signaling and affect surrounding cells' function and behavior. Cytokines include interferons, tumor necrosis factors, lymphokines, chemokines, and interleukins secreted by immune cells, mast cells, and various stromal cells (Akhter, Wu, Memon, & Mohsin, 2015).

Chemokines are a family of cytokines involved in the direct migration of leukocytes and activation of inflammatory stimuli. Chemokines and their receptors play a vital role in the growth, survival, or death of tumor cells as well as their metastasis

(Jorgensen et al., 2019; Moore, 2001). Chemokine C-C motif ligand 21 (CCL21) is a cytokine that binds specifically to the CCR7 chemokine receptor, which has antitumor properties and can predict tumorigenesis in cancer (Madej et al., 2013; McHugh, 2019). CCL21/CCR7 has essential roles in immune cell and lymph-node homing, peripheral tolerance, development and function of T regulatory cells, and lymphoid neogenesis (Joutoku et al., 2019; Zhao et al., 2014). Increased CD8+ T cells can reduce the progression of viral diseases such as HIV and COVID-19. High expression of these genes acts as biomarkers in various diseases such as cancers and viral infections like HIV infection and pneumonia (Cyster, 1999; Gollmer et al., 2009; Jorgensen et al., 2019). CCL21 is known as a base for cancer immunotherapy since it can chemoattract T lymphocytes and DCs (Beemiller, Jacobelli, & Krummel, 2012). DCs receive tumor antigens and migrate to T-cell zones of lymphoid organs for particular antitumor T-cell activity (Zhao et al., 2014). Interleukin-1 beta (IL1 β) belongs to the family of cytokines with severe proinflammation and, the *IL1 β* gene encodes it in humans. IL1 β is involved in various cellular

* Corresponding author's e-mail address:

maria_beihaghi@yahoo.com

[†]These authors contributed equally to this work.

activities such as activation of neutrophils, T and B lymphocytes, production of cytokines, antibodies, collagen, and fibroblast proliferation. (Van Damme et al., 1985). VQGEESNDK epitope is a part of IL-1 β that acts as a potent adjuvant by binding to secretory protein sequences. This peptide sequence possesses all of the IL-1 β adjuvant activity without any inflammatory response, such as induction of a fever response. (Boraschi, Tagliabue, & Miller, 2009).

Molecular dynamic simulation (MD) is a computer simulation method used to understand the conformational changes in recombinant proteins due to mutations comparatively (Musiani et al., 2014). In addition, its dynamics information can be used to analyze the highly fluctuating and complex nature of protein dynamics (Gaieb & Morikis, 2017).

The project's goal was to use immunoinformatic methods based on drug design algorithms to simulate and produce a drug based upon cytokine genes. Therefore, due to the complexity of the discovery process and the efficiency of neural network techniques, in addition to molecular docking, neural network techniques with neurophase rules were used to design an efficient diagnostic model in the immune system.

Materials and Methods

Construction of Amino acid Cassette

The epitope of human CCL21 (Accession No: CAG29322.1) and epitope of human IL-1 β (PDB ID: 4G6M_A) were designed as the principal part of the recombinant protein cassette. So, the best T cell epitope of CCL21, EAAAK sequence as beta-defensins linker, the epitope of human IL-1 beta, RRVR as sensitive foreign protease linker, the signal SEKDEL for effectual agglomeration of the plant recombinant protein in the endoplasmic reticulum (ER), RVLAEA sequence as HIV protease linker and 6xHis tags was possessed (Figure 1)

Prediction of T- and B-Cell Epitopes of the Recombinant Protein

T-cell and B-cell epitopes of this recombinant protein were identified by BepiPred 2.0, BCpreds, ABCpreds, SVMTrip, and MAPPP online servers (Table 1 and 2). These bioinformatic tools have been used to prognosticate antigenic epitopes presented on the T-cell and B-cell surface by major histocompatibility complex class I and II molecules (MHC I, MHCII) (O'Donnell, Rubinsteyn, & Laserson, 2020). TAPPred server was used to verify the recombinant protein binding affinity of peptides toward the TAP transporter. This online service is based on cascade SVM and uses the amino acids sequence and properties.




Figure 1. A) Schematic representation of CCL21/IL1 β protein construct.

Table 1. Prediction of T- cell epitopes of human CCL21 (PDB ID: 2L4N). A covering score over 90% and IC50 below 50 were the best binder epitopes

Epitope peptide	MHCI		Epitope peptide	MHCII	
	allele HLA	score		allele HLA	IC50
IPAKVVRSY	HLA-B*35:01	99.4%	LWVQQLMQH	HLA-DRB4*01:01	15.60
LPRKRSQAEL	HLA-B*07:02	95.7%	ELWVQQLMQ	HLA-DRB4*01:01	17.10
LCADPKELW	HLA-B*58:01	95.4%	PKELWVQQ	HLA-DRB4*01:01	18.50
LCADPKELW	HLA-B*57:01	93.4%	AKVVRSYRK	HLA-DRB4*01:01	19.20
IPAKVVRSY	HLA-B*53:01	90.3%	QQLMQHLDK	HLA-DRB4*01:01	21.40

Table 2. Prediction of T-cell epitopes of CCL21/IL1 β protein.

MHCI			MHCII		
Epitope peptide	allele HLA	score	Epitope peptide	allele HLA	IC50
IPAKVVRSY	HLA-B*35:01	99.4%	SGTND AEDCCLSVTQ	HLA-DRB1*08:02, HLA-DRB5*01	14.20
KELWVQQLM	HLA-B*40:01	93%	CAPPDQPWVERIIQR	HLA-DRB1*04:01	16.20

A covering score over 90% w and IC50 below 50 were the best binder epitopes of MHC I and MHC II-related HLAs, respectively.

Prediction of Physicochemical Characterization of Recombinant Protein

The SOLpro server measured the solubility of recombinant protein. Furthermore, ProtParam online server was used to identify various physicochemical parameters, including amino acid composition, pI, aliphatic index (II), instability index, in vivo and in vitro half-life, molecular weight (MW), and grand average of hydropathicity (GRAVY). The SignalP 5.0 server predicts the presence of signal peptides and the location of their cleavage sites in recombinant and commercial proteins. Localization of protein was analyzed by DeepLoc online server. The allergenicity of this recombinant protein was assessed by the AllgPred web server, which was used to show the post-translational modifications of CCL21/IL1 β . NetOGlyc 4.0 Server was used to show the O-glycosylation and NetNGlyc 1.0 Server to show the N-glycosylation site of this recombinant protein. NetPhos 3.1 Server was used to find the phosphorylation sites of the protein (Safavi et al., 2019).

Molecular Dynamic Simulation and the Prediction of the Stability and Flexibility of the Recombinant Protein

PSIPRED webserver was used for computational modeling and getting the PDB file of the CCL21/IL-1 β construct. Molecular dynamic (MD) simulations were utilized using GROMACS-4.5 and GROMOS96 (ffG45a3) force fields to assess the conformational changes of the protein. In order to neutralize the system in terms of charge, 5 counter Cl⁻ ions for CCL21 and 7 counter Cl⁻ ions for CCL21/IL-1 β simulation were added to the solvated system. Pressure and temperature were kept at 1 bar and 300 K, and the system ran for 20 nanoseconds. The root-mean-square deviation (RMSD), the root-mean-square fluctuation (RMSF), and the radius of gyration of C-alpha atoms were calculated and analyzed using the Grace software. Also, the PDB files of the two proteins were aligned using the Pymol software.

Homology Modeling Structure Using *in silico* Tools

Since the 3D structure of this recombinant protein is not available, the comparative modeling method was used to create its three-dimensional




Figure 2. Prediction of the presence of signal peptides in commercial CCL21 and CCL21/IL1β recombinant protein by the SignalP 5.0 server; A) presence of signal peptides in 23 primary amino acids the location of their cleavage sites in 24th amino acid of commercial CCL21 were detected. B) as shown in this figure, no signal peptides in 65 primary amino acids of CCL21/IL1β recombinant protein.




Figure 3. Schematic representation of subcellular localizations CCL21/IL1β protein construct. This soluble and extracellular protein that localized in the nucleus with a Likelihood of 89.9 percent.

Table 3. Prediction of subcellular localization of CCL21/ IL1β constructs in human cells.

Localization	Nucleus	Cytoplasm	Cell membrane	Extracellular	Mitochondrion	Endoplasmic reticulum	Golgi apparatus	Plastid	Peroxisome	Lysosome/Vacuole
Likelihood	0.8918	0.093	0.015	0.0079	0.0009	0.0005	0.0005	0.002	0.0001	0
Type	Soluble	Membrane								
Likelihood	0.8434	0.1566								

Prediction of T- cell and B-cell Epitopes of the Recombinant Protein

T-cell and B-cell epitopes of recombinant protein were identified. All T cell epitopes of human CCL21 and CCL21/IL1β were illustrated in

Tables 1 and 2. There were no B cell epitopes predicted in human CCL21, but the VQGESNDK sequence of IL1 β was predicted as a B cell epitope.


Protein Structure Conformational Flexibility and Stability Analysis

In this project, we used MD simulations to compare the conformational changes of native and recombinant proteins. Multiple were analyzed throughout the simulation project, chiefly root mean square deviation (RMSD), root mean square fluctuations (RMSF), and the radius of gyration of the proteins with the time-dependent function of

MD. Obtained results proved that the dynamic motions of the two proteins are very similar.

RMSD values of CC121 commercial antigen, native protein, CCL21/IL-1 β recombinant antigen, and mutant proteins were analyzed to identify the effect of mutations on recombinant protein structure. We calculated RMSD for protein backbones and found RMSD values from the mutant structures to be quite stable, like the native protein.

The CC121 antigen and recombinant protein were stabilized at an RMSD value of around 5 Å, demonstrating that the mutations did not destabilize the protein structure (Figure 4A).






Figure 4. Protein structure conformational flexibility and stability analysis of CCI21 constructs, as the recombinant protein, and commercial CCI21, as the native protein, for 20 ns MD simulations. A) RMSD values during MD simulations of the recombinant protein and native protein structure. CCI21 antigen and recombinant protein were stabilized at an RMSD value of around 5 Å. B) Calculated average RMSF for C α atoms of the recombinant and native protein structure, residues located between Positions 30 and 60, residue fluctuations for CCL21/IL-1 β were similar to CCI21 commercial antigen and fairly low C) Radius of Gyration for the recombinant and native protein. The results of the radius of gyration indicated that CCI21/IL-1 β and CCI21 commercial antigen have the minimum compactness of their structures with 11.8 Å value. E) Visualization of the native (blue) and mutant (red) PDB files, aligned with PyMol software to show the same structure and conformational characteristics of these two proteins. F) The molecular docking of the complex of CCR7 and CCI21/IL-1 β . G) The molecular docking of the complex of CCR7 and human CCI21.

We also analyzed the RMSF fluctuations of each residue to specify the effect of mutations on protein residues. As Figure 4B demonstrates for residues located between positions 30 and 60, residue fluctuations for CCL21/IL-1 β were similar to CCI21 commercial antigen and fairly low.

The results of the radius of gyration indicated that CCI21/IL-1 β and CCI21 commercial antigen have the minimum compactness of their structures with 11.8 Å value. These data show that CCI21 mutations did not cause structural destabilizing effects, and no significant alterations were found for either protein's compactness during the simulation (Figure 4C).

Visualization Analysis of Native and Recombinant Protein

<http://jcmr.um.ac.ir>

The PDB files of two proteins were aligned together by PyMol software to compare the conformational differences of the native and recombinant proteins (Figure 4E). Put together. Our findings verify that these proteins have the same structural and conformational characteristics.

Comparative Analysis of Molecular Docking of these Two Proteins with CCR7 Receptor

This project aimed to investigate and comparative analysis of interactions between these proteins and CCR7 receptors. As shown in Table 3, molecular docking of human CCL21 was determined, and the best cluster had a score of -27, with a size of 34 complexes. In addition, Z-score was equal to -2.3. Molecular docking of CCI21/IL-1 β was also determined. The score of the best

Table 4. The best cluster of molecular docking result

protein	HADDOCK score	Cluster size	Z-Score
human CCL21	-27	34	-2.3
CCL21/IL1 β	-30	38	-2.8

cluster was -30, with a size of 38 complexes. In addition, Z-score was equal to -2.8 (Table 4). Results obtained from molecular docking of these two proteins were so similar, and as mentioned before, there were interactions between CCL21/IL1 β ligands and CCR7 receptors as human CCL21. The image of the complex of CCR7 - CCL21/IL-1 β and CCR7 - CCL21 has been shown in Figures 4 F & 4 G.

Conclusion

The findings of this experiment confirmed that the recombinant protein and commercial CCL21 have the same structural and conformational characteristics. Therefore, this recombinant protein maybe has the same function as commercial CCL21, like anti-metastatic and cytotoxicity effects on cancer cell lines. Also, it has a chemotactic response on lymphocyte cells and is a potential treatment option in cancer immunotherapy. We have achieved our predetermined goals: to produce the recombinant protein in different expression hosts like yeast and plant to improve the production of CCL21 recombinant protein. Other diagnostic tests should be performed before the clinical application and commercialization of this protein.

Acknowledgments

This study was financially supported by the Biotechnology Development Council of the Islamic Republic of Iran. We would also like to thank faculty of science, Ferdowsi university of Mashhad, Mashhad, Iran.

References

Akhter, N., Wu, B., Memon, A. M., and Mohsin, M. (2015). Probiotics and prebiotics associated with aquaculture: a review. *Fish & shellfish immunology*, 45(2), 733-741 .

Beemiller, P., Jacobelli, J., and Krummel, M. F. (2012). Integration of the movement of signaling microclusters with cellular motility in immunological synapses. *Nature immunology*, 13(8), 787-795 .

Boraschi, D., Tagliabue, A., and Miller, A. D. (2009). The immunostimulatory effect of IL-1 β in vivo is blocked by antisense peptides complementary to the loop sequence 163–171. *FEBS letters*, 583(4), 792-796 .

Cyster, J. G. (1999). Chemokines and the homing of dendritic cells to the T cell areas of lymphoid organs. *Journal of Experimental Medicine*, 189(3), 447-450 .

Gaieb, Z., & Morikis, D. (2017). Detection of Side Chain Rearrangements Mediating the Motions of Transmembrane Helices in Molecular Dynamics Simulations of G Protein-Coupled Receptors. *Computational and structural biotechnology journal*, 15, 131-137 .

Gollmer, K., Asperti-Boursin, F., Tanaka, Y., Okkenhaug, K., Vanhaesebroeck, B., Peterson, J. R., et al. (2009). CCL21 mediates CD4+ T-cell costimulation via a DOCK2/Rac-dependent pathway. *Blood*, 114(3), 580-588 .

Jorgensen, A. S., Rosenkilde, M. M., Hjortø, G. M., Larsen, O., Legler, D. F., Uetz-von Allmen, E., et al. (2019). Biased signaling of CCL21 and CCL19 does not rely on N-terminal differences, but markedly on the chemokine core domains and extracellular loop 2 of CCR7. *Frontiers in immunology*, 10, 2156 .

Joutoku, Z., Onodera, T., Matsuoka, M., Homan, K., Momma, D., Baba, R., Hishimura, R. (2019). CCL21/CCR7 axis regulating juvenile cartilage repair can enhance cartilage healing in adults. *Scientific reports*, 9(1), 5165 .

Madej, T., Lanczycki, C. J., Zhang, D., Thiessen, P. A., Geer, R. C., Marchler-Bauer, A., and Bryant, S. H. (2013). MMDB and VAST+: tracking structural similarities between macromolecular complexes. *Nucleic acids research*, 42(D1), D297-D303 .

McHugh, J. (2019). CCL21–CCR7 axis in RA: linking inflammation and bone erosion. *Nature Reviews Rheumatology*, 15(10), 576-576 .

Moore, M. A. (2001). The role of chemoattraction in cancer metastases. *Bioessays*, 23(8), 674-676 .

O'Donnell, T. J., Rubinsteyn, A., and Laserson, U.

(2020). MHCflurry 2.0: Improved Pan-Allele Prediction of MHC Class I-Presented Peptides by Incorporating Antigen Processing. *Cell Systems*, 11(1), 42-48. e47 .

Van Damme, J., De Ley ,M., Opdenakker, G., Billiau, A., De Somer, P., and Van Beeumen, J. (1985). Homogeneous interferon-inducing 22K factor is related to endogenous pyrogen and interleukin-1. *Nature*, 314(6008), 266-268 .

Zhao, D.-x., Li, Z.-j., Zhang, Y., Zhang, X.-n., Zhao, K.-c., Li, Y.-g., et al. (2014). Enhanced antitumor immunity is elicited by adenovirus-mediated gene transfer of CCL21 and IL-15 in murine colon carcinomas. *Cellular immunology*, 289(1), 155-161 .

Open Access Statement:

This is an open access article distributed under the Creative Commons Attribution License (CC-BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.